

Product datasheet for TL507326

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Prkag3 Mouse shRNA Plasmid (Locus ID 241113)

Product data:

Product Type: shRNA Plasmids

Product Name: Prkag3 Mouse shRNA Plasmid (Locus ID 241113)

Locus ID: 241113

Synonyms: Amkg3; Ampkg3; AMPKg3L; AMPKg3S

Vector:pGFP-C-shLenti (TR30023)E. coli Selection:Chloramphenicol (34 ug/ml)

Mammalian Cell

Puromycin

Selection:

Format: Lentiviral plasmids

Components: Prkag3 - Mouse, 4 unique 29mer shRNA constructs in lentiviral GFP vector(Gene ID = 241113).

5µg purified plasmid DNA per construct

29-mer scrambled shRNA cassette in pGFP-C-shLenti Vector, TR30021, included for free.

RefSeq: <u>BC116749</u>, <u>BC116777</u>, <u>NM 153744</u>, <u>NM 153745</u>, <u>NM 153744.1</u>, <u>NM 153744.2</u>, <u>NM 153744.3</u>

UniProt ID: Q8BGM7

Summary: AMP/ATP-binding subunit of AMP-activated protein kinase (AMPK), an energy sensor protein

kinase that plays a key role in regulating cellular energy metabolism. In response to reduction of intracellular ATP levels, AMPK activates energy-producing pathways and inhibits energy-consuming processes: inhibits protein, carbohydrate and lipid biosynthesis, as well as cell growth and proliferation. AMPK acts via direct phosphorylation of metabolic enzymes, and by longer-term effects via phosphorylation of transcription regulators. Also acts as a regulator of cellular polarity by remodeling the actin cytoskeleton; probably by indirectly activating myosin. Gamma non-catalytic subunit mediates binding to AMP, ADP and ATP, leading to activate or inhibit AMPK: AMP-binding results in allosteric activation of alpha catalytic subunit (PRKAA1 or PRKAA2) both by inducing phosphorylation and preventing dephosphorylation of catalytic subunits. ADP also stimulates phosphorylation, without stimulating already phosphorylated catalytic subunit. ATP promotes dephosphorylation of catalytic subunit,

shRNA Design: These shRNA constructs were designed against multiple splice variants at this gene locus. To

be certain that your variant of interest is targeted, please contact <u>techsupport@origene.com</u>. If you need a special design or shRNA sequence, please utilize our <u>custom shRNA service</u>.

rendering the AMPK enzyme inactive (By similarity).[UniProtKB/Swiss-Prot Function]







Performance Guaranteed:

OriGene guarantees that the sequences in the shRNA expression cassettes are verified to correspond to the target gene with 100% identity. One of the four constructs at minimum are guaranteed to produce 70% or more gene expression knock-down provided a minimum transfection efficiency of 80% is achieved. Western Blot data is recommended over qPCR to evaluate the silencing effect of the shRNA constructs 72 hrs post transfection. To properly assess knockdown, the gene expression level from the included scramble control vector must be used in comparison with the target-specific shRNA transfected samples.

For non-conforming shRNA, requests for replacement product must be made within ninety (90) days from the date of delivery of the shRNA kit. To arrange for a free replacement with newly designed constructs, please contact Technical Services at techsupport@origene.com. Please provide your data indicating the transfection efficiency and measurement of gene expression knockdown compared to the scrambled shRNA control (Western Blot data preferred).