

## **Product datasheet for TP314681**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## HMGA2 (NM\_003484) Human Recombinant Protein

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant protein of human high mobility group AT-hook 2 (HMGA2), transcript variant 2, 20

μ

Species: Human Expression Host: HEK293T

Expression cDNA >RC214681 representing NM\_003484
Clone or AA Red=Cloning site Green=Tags(s)

Sequence:

MSARGEGAGQPSTSAQGQPAAPAPQKRGRGRPRKQQQEPTGEPSPKRPRGRPKGSKNKSPSKAAQKKAEA

TGEKRPRGRPRKWDNLLPRTSSKKKTSLGNSTKRSH

**TRTRPLEQKLISEEDLAANDILDYKDDDDKV** 

Tag: C-Myc/DDK

**Predicted MW:** 11.3 kDa

**Concentration:** >0.05 μg/μL as determined by microplate BCA method

**Purity:** > 80% as determined by SDS-PAGE and Coomassie blue staining

**Buffer:** 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol

**Preparation:** Recombinant protein was captured through anti-DDK affinity column followed by conventional

chromatography steps.

**Note:** For testing in cell culture applications, please filter before use. Note that you may experience

some loss of protein during the filtration process.

Storage: Store at -80°C.

Stability: Stable for 12 months from the date of receipt of the product under proper storage and handling

conditions. Avoid repeated freeze-thaw cycles.

**RefSeq:** NP 003475

 Locus ID:
 8091

 UniProt ID:
 P52926

 RefSeq Size:
 1539



Cytogenetics: 12q14.3

RefSeq ORF: 318

Synonyms: BABL; HMGI-C; HMGIC; LIPO; SRS5; STQTL9

Summary: This gene encodes a protein that belongs to the non-histone chromosomal high mobility group

(HMG) protein family. HMG proteins function as architectural factors and are essential


components of the enhancesome. This protein contains structural DNA-binding domains and may act as a transcriptional regulating factor. Identification of the deletion, amplification, and rearrangement of this gene that are associated with myxoid liposarcoma suggests a role in adipogenesis and mesenchymal differentiation. A gene knock out study of the mouse counterpart demonstrated that this gene is involved in diet-induced obesity. Alternate

transcriptional splice variants, encoding different isoforms, have been characterized. [provided

by RefSeq, Jul 2008]

**Protein Families:** Druggable Genome

## **Product images:**



Coomassie blue staining of purified HMGA2 protein (Cat# TP314681). The protein was produced from HEK293T cells transfected with HMGA2 cDNA clone (Cat# [RC214681]) using MegaTran 2.0 (Cat# [TT210002]).