

## Product datasheet for TP319216

### C4BPB (NM\_001017367) Human Recombinant Protein

#### Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Homo sapiens complement component 4 binding protein, beta (C4BPB), transcript variant 5, 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC219216 protein sequence <b>Red</b> =Cloning site <b>Green</b> =Tags(s)
	 MFFWCACCLMVAWRVSASDEHCPPELPPVDNSIFVAKEVEGQILGTYVCIKGYHLVGKKTLCFNASKEDWN TTTECRLGHCPDPVLVNGEFSSSGPVNVSDKITFMCNDHYILKGSNRSQCLEDHTWAPPFPICKSRDCDP PGNPVHGYFEGNNFTLGSTISYYCEDRYLVGVQEQQCVDGEWSSALPVCKLIQEAPKPECEKALLAFQE SKNLCEAMENFMQQLKESGMTMEELKYSLELKKALAKLL  <b>TRTRPLEQKLISEEDLAANDILDYKDDDDKV</b>
Tag:	C-Myc/DDK
Predicted MW:	26.3 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
RefSeq:	<a href="#">NP_001017367</a>
Locus ID:	725




[View online »](#)

UniProt ID: [P20851](#)  
RefSeq Size: 967  
Cytogenetics: 1q32.1  
RefSeq ORF: 812  
Synonyms: C4BP

**Summary:** This gene encodes a member of a superfamily of proteins composed predominantly of tandemly arrayed short consensus repeats of approximately 60 amino acids. A single, unique beta-chain encoded by this gene assembles with seven identical alpha-chains into the predominant isoform of C4b-binding protein, a multimeric protein that controls activation of the complement cascade through the classical pathway. C4b-binding protein has a regulatory role in the coagulation system also, mediated through the beta-chain binding of protein S, a vitamin K-dependent protein that serves as a cofactor of activated protein C. The genes encoding both alpha and beta chains are located adjacent to each other on human chromosome 1 in the regulator of complement activation gene cluster. Alternative splicing gives rise to multiple transcript variants. [provided by RefSeq, Jul 2008]

**Protein Pathways:** Complement and coagulation cascades

### Product images:



Coomassie blue staining of purified C4BPB protein (Cat# TP319216). The protein was produced from HEK293T cells transfected with C4BPB cDNA clone (Cat# [RC219216]) using MegaTran 2.0 (Cat# [TT210002]).