

Product datasheet for **TP319802**

Sterol carrier protein 2 (SCP2) (NM_002979) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Recombinant protein of human sterol carrier protein 2 (SCP2), transcript variant 1, 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC219802 representing NM_002979 Red =Cloning site Green =Tags(s)

MSSSPWEPATLRRVVFVGVGMFKVFKPGAENSRDYPDLAEAGKKALADAQIPYSAVDQACVGYVFGDST
CGQRAIYHSLGMTGIPIINVNNNCATGSTALLFMARQLIQGGVAEVLALGFEEKMSKGLGIKFSDRTIPT
DKHVDLLINKYGLSAHPVAPQMFYAGKEHMEKYGTKIEHFAKIGWKNHKHSVNNPYSQFQDEYSLDEVM
ASKEVDFLTLQCCPTSDGAAAAILASEAFVQKYGLQSKAVEILAQEMMTDLPSSFEEKSIKMGVDFDM
SKEAARKCYEKSGLTPNDIDVIELHDCFSTNELLTYEALGLCEPQGATLVDRGDNTYGGKWINPSGGL
ISKGHPLGATGLAQCAELCWQLRGEAGKRQVPKAKVALQHNLGIGGAVVVTLYKMGFPEAASSFRTHQIE
AVPTSSASDGFKANLVFKEIEKKLEEEGEQFVKKIGGIFAFKVKDGPGGKEATWVVDVKNKGKSVLPNSD
KKADCTITMADSDFLALMTGKMNPQSAFFQGKLIKITGNMGLAMKLQNLQLQPGNAKL

TRTRPLEQKLISEEDLAANDILDYKDDDDKV

Tag:	C-Myc/DDK
Predicted MW:	58.8 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.



[View online >](#)

RefSeq: [NP_002970](#)

Locus ID: 6342

UniProt ID: [P22307](#), [A0A384NY87](#), [B2R761](#), [Q59HG9](#)

RefSeq Size: 2697

Cytogenetics: 1p32.3


RefSeq ORF: 1641

Synonyms: NLTP; NSL-TP; SCOX; SCP-2; SCP-CHI; SCP-X; SCPX

Summary: This gene encodes two proteins: sterol carrier protein X (SCPx) and sterol carrier protein 2 (SCP2), as a result of transcription initiation from 2 independently regulated promoters. The transcript initiated from the proximal promoter encodes the longer SCPx protein, and the transcript initiated from the distal promoter encodes the shorter SCP2 protein, with the 2 proteins sharing a common C-terminus. Evidence suggests that the SCPx protein is a peroxisome-associated thiolase that is involved in the oxidation of branched chain fatty acids, while the SCP2 protein is thought to be an intracellular lipid transfer protein. This gene is highly expressed in organs involved in lipid metabolism, and may play a role in Zellweger syndrome, in which cells are deficient in peroxisomes and have impaired bile acid synthesis. Alternative splicing of this gene produces multiple transcript variants, some encoding different isoforms.[provided by RefSeq, Aug 2010]

Protein Pathways: Metabolic pathways, PPAR signaling pathway, Primary bile acid biosynthesis

Product images:



Coomassie blue staining of purified SCP2 protein (Cat# TP319802). The protein was produced from HEK293T cells transfected with SCP2 cDNA clone (Cat# [RC219802]) using MegaTran 2.0 (Cat# [TT210002]).