

Product datasheet for TP320522

PML Protein (PML) (NM_033246) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Homo sapiens promyelocytic leukemia (PML), transcript variant 7, 20 µg
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	>RC220522 representing NM_033246 Red =Cloning site Green =Tags(s)
	MEPAPARSPRPQQDPARPQEPTMPPPETPSEGRQPSPSPSPTERAPASEEEFQFLRCQQCQAEAKCPKLL PCLHTLCSGCLEASGMQCPCQAPWPLGADTPALDNVFFESLQRRLSVYRQIVDAQAVCTRCKESADFWC FECEQLLCAKCFEAHQWFLKHEARPLAELRNQSVREFLDGTRKTNNIFCSNPNHRTPTLTSIYCRGCSKP LCCSCALLDSSHSELKCDISAEIQQRQEELDAMTQALQEQDSAFGAVHAQMHAAVGQLGRARAETEELIR ERVRQVVAHVRAQERELLEAVDARYQRDYEEASRLGRLDAVLQRIRTGSALVQRMKCYASDQEVLDMHG FLRQALCRLRQEEPQSLQAAVRTDGFDEFKVRQLQLSSCITQGKDAAVSKKASPEAASTPRDPIDVDLRN ALW
	SGP TRRRLEQKLISEEDLAANDILDYKDDDDKV
Tag:	C-Myc/DDK
Predicted MW:	47.4 kDa
Concentration:	>0.05 µg/µL as determined by microplate BCA method
Purity:	> 80% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol
Preparation:	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Note:	For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process.
Storage:	Store at -80°C.
Stability:	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.



[View online »](#)

RefSeq: [NP_150249](#)

Locus ID: 5371

UniProt ID: [P29590](#)

RefSeq Size: 1851

Cytogenetics: 15q24.1

RefSeq ORF: 1269


Synonyms: MYL; PP8675; RNF71; TRIM19

Summary: The protein encoded by this gene is a member of the tripartite motif (TRIM) family. The TRIM motif includes three zinc-binding domains, a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. This phosphoprotein localizes to nuclear bodies where it functions as a transcription factor and tumor suppressor. Its expression is cell-cycle related and it regulates the p53 response to oncogenic signals. The gene is often involved in the translocation with the retinoic acid receptor alpha gene associated with acute promyelocytic leukemia (APL). Extensive alternative splicing of this gene results in several variations of the protein's central and C-terminal regions; all variants encode the same N-terminus. Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Jul 2008]

Protein Families: Druggable Genome, Transcription Factors

Protein Pathways: Acute myeloid leukemia, Pathways in cancer, Ubiquitin mediated proteolysis

Product images:



Coomassie blue staining of purified PML protein (Cat# TP320522). The protein was produced from HEK293T cells transfected with PML cDNA clone (Cat# [RC220522]) using MegaTran 2.0 (Cat# [TT210002]).