

Product datasheet for **TP500154**

Snapap (BC006744) Mouse Recombinant Protein

Product data:

| | |
|---------------------------------------|---|
| Product Type: | Recombinant Proteins |
| Description: | Purified recombinant protein of Mouse SNAP-associated protein (cDNA clone MGC:12076 IMAGE:3708620), complete cds, with C-terminal MYC/DDK tag, expressed in HEK293T cells, 20ug |
| Species: | Mouse |
| Expression Host: | HEK293T |
| Expression cDNA Clone or AA Sequence: | >MR200154 protein sequence Red =Cloning site Green =Tags(s) |
| | MLVAHFLFPELCRINEDQKVALDLDPYVKLLNARRRVVLVNNILQNAQERLRRLNHSVAKETARRRAML DSGVYPPGSPSK |
| | TRTRPLEQKLISEEDLAANDILDYKDDDDKV |
| Tag: | C-MYC/DDK |
| Predicted MW: | 9.5 kDa |
| Concentration: | >0.05 µg/µL as determined by microplate BCA method |
| Purity: | > 80% as determined by SDS-PAGE and Coomassie blue staining |
| Buffer: | 25 mM Tris-HCl, 100 mM glycine, pH 7.3, 10% glycerol |
| Note: | For testing in cell culture applications, please filter before use. Note that you may experience some loss of protein during the filtration process. |
| Storage: | Store at -80°C after receiving vials. |
| Stability: | Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles. |
| Locus ID: | 20615 |
| UniProt ID: | Q9Z266 |
| RefSeq Size: | 1858 |
| Cytogenetics: | 3 F1 |
| RefSeq ORF: | 246 |



[View online »](#)

Synonyms: 25kDa; AA407989; AV026596; Bloc1s7; Snap25bp; Snapap

Summary: Component of the BLOC-1 complex, a complex that is required for normal biogenesis of lysosome-related organelles (LRO), such as platelet dense granules and melanosomes. In concert with the AP-3 complex, the BLOC-1 complex is required to target membrane protein cargos into vesicles assembled at cell bodies for delivery into neurites and nerve terminals. The BLOC-1 complex, in association with SNARE proteins, is also proposed to be involved in neurite extension. Plays a role in intracellular vesicle trafficking and synaptic vesicle recycling. May modulate a step between vesicle priming, fusion and calcium-dependent neurotransmitter release through its ability to potentiate the interaction of synaptotagmin with the SNAREs and the plasma-membrane-associated protein SNAP25. Its phosphorylation state influences exocytotic protein interactions and may regulate synaptic vesicle exocytosis. May also have a role in the mechanisms of SNARE-mediated membrane fusion in non-neuronal cells (PubMed:16760431, PubMed:19546860, PubMed:21998198). As part of the BORC complex may play a role in lysosomes movement and localization at the cell periphery. Associated with the cytosolic face of lysosomes, the BORC complex may recruit ARL8B and couple lysosomes to microtubule plus-end-directed kinesin motor (By similarity). [UniProtKB/Swiss-Prot Function]