

Product datasheet for TP720467M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CD8B (NM 001178100) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Recombinant protein of human CD8b molecule (CD8B), transcript variant 6.

Species: Human
Expression Host: HEK293

Expression cDNA Clone

Leu22-Pro170

or AA Sequence:

Tag: C-His

Predicted MW: 17.8 kDa

Concentration: lot specific

Purity: >95% as determined by SDS-PAGE and Coomassie blue staining

Buffer: Provided lyophilized from a 0.2 μm filtered solution of 20 mM Tris-HCl, 150 mM NaCl

Endotoxin: < 0.1 EU per μg protein as determined by LAL test

Reconstitution Method: Always centrifuge tubes before opening. Do not mix by vortex or pipetting. Dissolve the

lyophilized protein in ddH2O. It is not recommended to reconstitute a concentration less than 100 µg/ml. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

Storage: Store at -80°C.

Stability: Stable for at least 6 months from date of receipt under proper storage and handling

conditions.

RefSeq: NP 001171571

 Locus ID:
 926

 UniProt ID:
 P10966

 Cytogenetics:
 2p11.2

Synonyms: CD8B1; LEU2; LY3; LYT3; P37





Summary:

The CD8 antigen is a cell surface glycoprotein found on most cytotoxic T lymphocytes that mediates efficient cell-cell interactions within the immune system. The CD8 antigen, acting as a coreceptor, and the T-cell receptor on the T lymphocyte recognize antigens displayed by an antigen presenting cell (APC) in the context of class I MHC molecules. The functional coreceptor is either a homodimer composed of two alpha chains, or a heterodimer composed of one alpha and one beta chain. Both alpha and beta chains share significant homology to immunoglobulin variable light chains. This gene encodes the CD8 beta chain isoforms. Multiple alternatively spliced transcript variants encoding distinct membrane associated or secreted isoforms have been described. A pseudogene, also located on chromosome 2, has been identified. [provided by RefSeq, May 2010]

Protein Families:

Druggable Genome, Secreted Protein, Transmembrane

Protein Pathways:

Antigen processing and presentation, Cell adhesion molecules (CAMs), Hematopoietic cell lineage, Primary immunodeficiency, T cell receptor signaling pathway

Product images:

