

Product datasheet for **TP723292**

MDC (CCL22) (NM_002990) Human Recombinant Protein

Product data:

Product Type:	Recombinant Proteins
Description:	Purified recombinant protein of Human chemokine (C-C motif) ligand 22 (CCL22).
Species:	Human
Expression Host:	E. coli
Expression cDNA Clone or AA Sequence:	GPYGANMEDS VCCRDYVRYR LPLRWVKHFY WTSDSCPRPG VLLTFRDKE ICADPRVPWW KMILNKLSQ
Tag:	Tag Free
Predicted MW:	8.1 kDa
Concentration:	lot specific
Purity:	>95% as determined by SDS-PAGE and Coomassie blue staining
Buffer:	Lyophilized from a 0.2 µM filtered solution of 20mM phosphate buffer, 100mM NaCl, pH 7.2
Bioactivity:	Determined by its ability to chemoattract human T cells using a concentration range of 10.0-100.0 ng/ml.
Endotoxin:	Endotoxin level is < 0.1 ng/µg of protein (< 1 EU/µg)
Storage:	Store at -80°C.
Stability:	Stable for at least 6 months from date of receipt under proper storage and handling conditions.
RefSeq:	NP_002981
Locus ID:	6367
UniProt ID:	O00626
RefSeq Size:	2933
Cytogenetics:	16q21
RefSeq ORF:	279
Synonyms:	A-152E5.1; ABCD-1; DC/B-CK; MDC; SCYA22; STCP-1



[View online »](#)

Summary:

This antimicrobial gene is one of several Cys-Cys (CC) cytokine genes clustered on the q arm of chromosome 16. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for monocytes, dendritic cells, natural killer cells and for chronically activated T lymphocytes. It also displays a mild activity for primary activated T lymphocytes and has no chemoattractant activity for neutrophils, eosinophils and resting T lymphocytes. The product of this gene binds to chemokine receptor CCR4. This chemokine may play a role in the trafficking of activated T lymphocytes to inflammatory sites and other aspects of activated T lymphocyte physiology. [provided by RefSeq, Sep 2014]

Protein Families:

Druggable Genome, Secreted Protein, Transmembrane

Protein Pathways:

Chemokine signaling pathway, Cytokine-cytokine receptor interaction

Product images: