

## **Product datasheet for TP723455**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Thrombopoietin (THPO) (NM\_000460) Human Recombinant Protein

#### **Product data:**

**Product Type:** Recombinant Proteins

**Description:** Purified recombinant protein of Human thrombopoietin (THPO), transcript variant 1.

Species: Human
Expression Host: E. coli

**Expression cDNA Clone** 

or AA Sequence:

SPAPPACDLR VLSKLLRDSH VLHSRLSQCP EVHPLPTPVL LPAVDFSLGE WKTQMEETKA QDILGAVTLL LEGVMAARGQ LGPTCLSSLL GQLSGQVRLL LGALQSLLGT QLPPQGRTTA

HKDPNAIFLS FQHLLRGKVR FLMLVGGSTL CVRRAPPTTA VPSRTSLVLT LNEL

Tag: Tag Free
Predicted MW: 18.6 kDa
Concentration: lot specific

**Purity:** >95% as determined by SDS-PAGE and Coomassie blue staining

Buffer: Lyophilized from a 0.2 μM filtered solution of 20mM phosphate buffer,100mM NaCl, pH 7.2

**Bioactivity:** ED50 was determined by the dose-dependent stimulation of the proliferation of human

MO7e cells was found to be < 1.0 ng/ml, corresponding to a specific activity of > 1  $\times$  10<sup>6</sup>

units/mg.

**Endotoxin:** Endotoxin level is < 0.1 ng/µg of protein (< 1 EU/µg)

Storage: Store at -80°C.

**Stability:** Stable for at least 6 months from date of receipt under proper storage and handling

conditions.

**RefSeq:** <u>NP 000451</u>

 Locus ID:
 7066

 UniProt ID:
 P40225

 RefSeq Size:
 1805

 Cytogenetics:
 3q27.1

 RefSeq ORF:
 1059

Synonyms: MGDF; MKCSF; ML; MPLLG; THCYT1; TPO





**Summary:** 

Megakaryocytopoiesis is the cellular development process that leads to platelet production. The main functional protein encoded by this gene is a humoral growth factor that is necessary for megakaryocyte proliferation and maturation, as well as for thrombopoiesis. This protein is the ligand for MLP/C\_MPL, the product of myeloproliferative leukemia virus oncogene. Mutations in this gene are the cause of thrombocythemia 1. Alternative promoter usage and differential splicing result in multiple transcript variants differing in the 5' UTR and/or coding region. Multiple AUG codons upstream of the main open reading frame (ORF) have been identified, and these upstream AUGs inhibit translation of the main ORF at different extent. [provided by RefSeq, Feb 2014]

**Protein Families:** Druggable Genome, Secreted Protein

**Protein Pathways:** Hematopoietic cell lineage

### **Product images:**

