

Product datasheet for TP723713

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

IL6 (NM_000600) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Description: Purified recombinant protein of Human interleukin 6 (interferon, beta 2) (IL6)

Species: Human
Expression Host: E. coli

Expression cDNA Clone

or AA Sequence:

Human IL-6, the region of Pro29-Met212, from gene Accession# NM_000600

Tag: Tag Free
Predicted MW: 21 kDa
Concentration: lot specific

Purity: Purity is >98%, as determined by Coomassie stained SDS-PAGE.

Buffer: PBS, pH 7.2, 0.5M NaCl, 10% Glycerol

Bioactivity: The ED50 is 4-20 pg/ml, corresponding to a specific activity of 0.5-2.5 X10⁸ units/mg,

determined by a dose dependent stimulation in a 7TD1 cell proliferation assay.

Endotoxin: Less than 0.01 ng per ug protein as determined by the LAL method

Storage: Store at -80°C.

Stability: Unopened vial can be stored between 2°C and 8°C for up to 2 weeks, at -20°C for up to 6

months, or at -70°C or below until the expiration date. Aliquots can be stored between 2°C and 8°C for up to one week and stored at -20°C or colder for up to 3 months. Avoid repeated

freeze/thaw cycles.

RefSeq: NP 000591

Locus ID: 3569

UniProt ID: <u>P05231</u>, <u>Q75MH2</u>, <u>B4DVM1</u>

RefSeq Size: 1201 Cytogenetics: 7p15.3 RefSeq ORF: 636

Synonyms: BSF-2; BSF2; CDF; HGF; HSF; IFN-beta-2; IFNB2; IL-6





Summary:

This gene encodes a cytokine that functions in inflammation and the maturation of B cells. In addition, the encoded protein has been shown to be an endogenous pyrogen capable of inducing fever in people with autoimmune diseases or infections. The protein is primarily produced at sites of acute and chronic inflammation, where it is secreted into the serum and induces a transcriptional inflammatory response through interleukin 6 receptor, alpha. The functioning of this gene is implicated in a wide variety of inflammation-associated disease states, including suspectibility to diabetes mellitus and systemic juvenile rheumatoid arthritis. Elevated levels of the encoded protein have been found in virus infections, including COVID-19 (disease caused by SARS-CoV-2). [provided by RefSeq, Aug 2020]

Protein Families:

Druggable Genome, Secreted Protein

Protein Pathways:

Cytokine-cytokine receptor interaction, Cytosolic DNA-sensing pathway, Graft-versus-host disease, Hematopoietic cell lineage, Hypertrophic cardiomyopathy (HCM), Jak-STAT signaling pathway, NOD-like receptor signaling pathway, Pathways in cancer, Prion diseases, Toll-like receptor signaling pathway

Product images:

