

## **Product datasheet for TP724204**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **Human GDF15 Protein, hFc Tag**

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Human GDF15 Protein, hFc Tag

**Expression Host:** HEK293

Tag: N-Human Fc

**Predicted MW:** The protein has a predicted molecular mass of 38.4 kDa after removal of the signal

peptide. The apparent molecular mass of hFc-GDF15 is approximately 35-55 kDa due to

glycosylation.

**Purity:** The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie

blue staining.

**Reconstitution Method:** Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8% trehalose is added as protectants

before lyophilization.

Storage: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended

for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).

Lyophilized proteins are shipped at ambient temperature.

Stability: 12 months from date of despatch

**Summary:** This gene encodes a secreted ligand of the TGF-beta (transforming growth factor-beta)

superfamily of proteins. Ligands of this family bind various TGF-beta receptors leading to

recruitment and activation of SMAD family transcription factors that regulate gene

expression. The encoded preproprotein is proteolytically processed to generate each subunit of the disulfide-linked homodimer. The protein is expressed in a broad range of cell types, acts as a pleiotropic cytokine and is involved in the stress response program of cells after cellular injury. Increased protein levels are associated with disease states such as tissue hypoxia, inflammation, acute injury and oxidative stress. [provided by RefSeq, Aug 2016]