

## **Product datasheet for TP727374**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **Edar Mouse Recombinant Protein**

**Product data:** 

**Product Type:** Recombinant Proteins

**Description:** Recombinant Mouse Ectodysplasin Receptor/EDAR (C-Fc)

Species: Mouse

**Expression cDNA Clone** 

or AA Sequence:

Glu27-lle189

Tag: C-Fc

**Buffer:** Lyophilized from a 0.2 um filtered solution of PBS, pH 7.4.

**Note:** Recombinant Mouse Ectodysplasin Receptor is produced by our Mammalian expression

system and the target gene encoding Glu27-Ile189 is expressed with a Fc tag at the C-

terminus.

**Stability:** 12 months from date of despatch

Locus ID: 13608 UniProt ID: Q9R187

**Summary:** Ectodysplasin A receptor (EDAR) is a type I transmembrane protein of the TNF-α receptor

superfamily which plays a key role in ectodermal differentiation. EDAR was encoded by the mouse downless gene and defective in human dominant and recessive forms of autosomal hypohidrotic ectodermal dysplasia (EDA) syndrome. The extracellular domain of EDAR contains 14 cysteine residues, six of which approximate the TNFRSF cysteine-rich region, the cytoplasmic domain contains a region with homology to the death domains found in other TNFRSF members. EDAR has been suggested to be an early and important promoter of placode development in all ectodermal organs, such as uch as hair follicles, teeth and sweat glands. EDA-A1, the A1 isoform of EDA, is the EDAR ligand. EDA and EDA are implicated in appendage development by the cloning of a gene underlying hypohidrotic ectodermal dysplasia (HED) in mouse and human. HED is characterized by agenesis or malformation of ectoderm-derived appendages, such as teeth, sweat glands and hair follicles, while the skin

itself develops normally.

