

Product datasheet for TP750006

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

IL6 (NM 000600) Human Recombinant Protein

Product data:

Product Type: Recombinant Proteins

Recombinant protein of human Interleukin 6(IL-6) produced in E. coli. Description:

Species: Human E. coli **Expression Host:**

Expression cDNA Clone

or AA Sequence:

A DNA sequence encoding the region (Pro29-Met212) of IL6

Tag: Tag Free Predicted MW: 20.9 kDa

Concentration: Resuspend the protein to the desired concentration in proper buffer.

Purity: >95% as determined by SDS-PAGE and Coomassie blue staining **Buffer:** Lyophilized from a sterile solution containing 20 mM PB, pH 7.4

Determined by the dose-dependant stimulation of murine 7TD1 cells. ED50 for this effect is **Bioactivity:**

typically 0.009-0.014 ng/ml.

Endotoxin: < 0.1 EU per 1 µg of the protein by the LAL

For testing in cell culture applications, please filter before use. Note that you may experience Note:

some loss of protein during the filtration process.

Store at -80°C. Storage:

Stability: Stable for 12 months from the date of receipt of the product under proper storage and

handling conditions. Avoid repeated freeze-thaw cycles.

RefSeq: NP 000591

Locus ID: 3569

UniProt ID: P05231, Q75MH2, B4DVM1

RefSeg Size: 1201

Cytogenetics: 7p15.3

RefSeq ORF: 636

Synonyms: BSF-2; BSF2; CDF; HGF; HSF; IFN-beta-2; IFNB2; IL-6





Summary:

This gene encodes a cytokine that functions in inflammation and the maturation of B cells. In addition, the encoded protein has been shown to be an endogenous pyrogen capable of inducing fever in people with autoimmune diseases or infections. The protein is primarily produced at sites of acute and chronic inflammation, where it is secreted into the serum and induces a transcriptional inflammatory response through interleukin 6 receptor, alpha. The functioning of this gene is implicated in a wide variety of inflammation-associated disease states, including suspectibility to diabetes mellitus and systemic juvenile rheumatoid arthritis. Elevated levels of the encoded protein have been found in virus infections, including COVID-19 (disease caused by SARS-CoV-2). [provided by RefSeq, Aug 2020]

Protein Families:

Druggable Genome, Secreted Protein

Protein Pathways:

Cytokine-cytokine receptor interaction, Cytosolic DNA-sensing pathway, Graft-versus-host disease, Hematopoietic cell lineage, Hypertrophic cardiomyopathy (HCM), Jak-STAT signaling pathway, NOD-like receptor signaling pathway, Pathways in cancer, Prion diseases, Toll-like receptor signaling pathway

Product images:

