

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TS424189P5

## Histone H1.5 (HIST1H1B) CytoSection


## Product data:

Expression Host:HEK293TExpression cDNA Clone or AA Sequence:TrueORF Clone RC224189Tag:C-MYC/DDKDetection Antibodies:DDK Rabbit monoclonal antibody, recognizing both N- and C-terminal tags (TA592569)ACCN:MM 005322, NP 005313Synonyms:H1; H1.5; H1B; H1F5; H1s-3; HIST1H1BStorage:Room TemperatureStability:Slides are guaranteed for a year from the date of receipt if proper storage instructions were followed.Preparation:HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cut and mounted on positively charged SuperFrost slides.	Product Type:	CytoSections
Expression Host:HEK293TExpression cDNA Clone or AA Sequence:TrueORF Clone RC224189Tag:C-MYC/DDKDetection Antibodies:DDK Rabbit monoclonal antibody, recognizing both N- and C-terminal tags (TA592569)ACCN:NM 005322, NP 005313Synonyms:H1; H1.5; H1B; H1F5; H1s-3; HIST1H1BStorage:Room TemperatureStability:Slides are guaranteed for a year from the date of receipt if proper storage instructions were followed.Preparation:HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cut and mounted on positively charged SuperFrost slides.	Description:	
Expression cDNA Clone or AA Sequence:TrueORF Clone RC224189Tag:C-MYC/DDKDetection Antibodies:DDK Rabbit monoclonal antibody, recognizing both N- and C-terminal tags (TA592569)ACCN:NM 005322, NP 005313Synonyms:H1; H1.5; H1B; H1F5; H1s-3; HIST1H1BStorage:Room TemperatureStability:Slides are guaranteed for a year from the date of receipt if proper storage instructions were followed.Preparation:HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cut and mounted on positively charged SuperFrost slides.	Species:	Human
or AA Sequence:Tag:C-MYC/DDKDetection Antibodies:DDK Rabbit monoclonal antibody, recognizing both N- and C-terminal tags (TA592569)ACCN:NM 005322, NP 005313Synonyms:H1; H1.5; H1B; H1F5; H1s-3; HIST1H1BStorage:Room TemperatureStability:Slides are guaranteed for a year from the date of receipt if proper storage instructions were followed.Preparation:HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cut and mounted on positively charged SuperFrost slides.	Expression Host:	HEK293T
Detection Antibodies:DDK Rabbit monoclonal antibody, recognizing both N- and C-terminal tags (TA592569)ACCN:NM 005322, NP 005313Synonyms:H1; H1.5; H1B; H1F5; H1s-3; HIST1H1BStorage:Room TemperatureStability:Slides are guaranteed for a year from the date of receipt if proper storage instructions were followed.Preparation:HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cut and mounted on positively charged SuperFrost slides.	•	TrueORF Clone RC224189
ACCN:NM 005322, NP 005313Synonyms:H1; H1.5; H1B; H1F5; H1s-3; HIST1H1BStorage:Room TemperatureStability:Slides are guaranteed for a year from the date of receipt if proper storage instructions were followed.Preparation:HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cut and mounted on positively charged SuperFrost slides.	Tag:	C-MYC/DDK
Synonyms:H1; H1.5; H1B; H1F5; H1s-3; HIST1H1BStorage:Room TemperatureStability:Slides are guaranteed for a year from the date of receipt if proper storage instructions were followed.Preparation:HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cut and mounted on positively charged SuperFrost slides.	Detection Antibodies:	DDK Rabbit monoclonal antibody, recognizing both N- and C-terminal tags (TA592569)
Storage:Room TemperatureStability:Slides are guaranteed for a year from the date of receipt if proper storage instructions were followed.Preparation:HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cut and mounted on positively charged SuperFrost slides.	ACCN:	<u>NM 005322, NP 005313</u>
Stability:Slides are guaranteed for a year from the date of receipt if proper storage instructions were followed.Preparation:HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cut and mounted on positively charged SuperFrost slides.	Synonyms:	H1; H1.5; H1B; H1F5; H1s-3; HIST1H1B
followed. Preparation: HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cut and mounted on positively charged SuperFrost slides.	Storage:	Room Temperature
were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cut and mounted on positively charged SuperFrost slides.	Stability:	
	Preparation:	were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cut
<b>Note:</b> This product is for research use only and is not approved for use in humans or in clinical diagnosis.	Note:	This product is for research use only and is not approved for use in humans or in clinical diagnosis.
<b>RefSeq:</b> <u>NP 005313</u>	RefSeq:	<u>NP 005313</u>
Locus ID: 3009	Locus ID:	3009
Cytogenetics: 6p22.1	Cytogenetics:	6p22.1




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### **Product images:**



Immunohistology staining on HIST1H1B overexpressed cytosection [TS424189] with mouse anti DDK clone OTI11C3 C/N [TA180144] at 1:400 dilution 4C O/N. Antibody staining was achieved with HIER Citrate pH6 , Polink1 with DAB chromogen detection kit ([D11-18]). Positive stain shown with the brown chromogen present. HEK293T cells were transfected with cDNA clone [RC224189], 5 micron sections, 40x magnification



Immunohistology staining on HIST1H1B overexpressed cytosection TS424189P5 with mouse anti DDK clone OTI11C3 C/N [TA180144] at 1:400 dilution 4C O/N. Antibody staining was achieved with HIER Citrate pH6 , Polink1 with DAB chromogen detection kit ([D11-18]). Positive stain shown with the brown chromogen present. HEK293T cells were transfected with cDNA clone [RC200548], 5 micron sections, 40x magnification

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US