

Product datasheet for **UM800032**

Proteasome subunit alpha type 6 (PSMA6) Mouse Monoclonal Antibody [Clone ID: UMAB102]

Product data:

Product Type:	Primary Antibodies
Clone Name:	UMAB102
Applications:	IHC, WB
Recommended Dilution:	IHC 1:100, IF 1:100
Reactivity:	Human, Mouse, Rat, Monkey, Dog
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human PSMA6 (NP_002782) produced in E.coli.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.5~1.0 mg/ml (Lot Dependent)
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	27.2 kDa
Gene Name:	proteasome 20S subunit alpha 6
Database Link:	NP_002782 Entrez Gene 26443 Mouse Entrez Gene 29673 Rat Entrez Gene 480290 Dog Entrez Gene 695113 Monkey Entrez Gene 5687 Human P60900



[View online »](#)

Background:

The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the peptidase T1A family, that is a 20S core alpha subunit. A pseudogene has been identified on the Y chromosome. [provided by RefSeq, Jul 2008]

Synonyms:


IOTA; p27K; PROS27

Protein Families:


Druggable Genome, Protease, Stem cell - Pluripotency

Protein Pathways:


Proteasome

Product images:


Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human breast tissue using anti-PSMA6 mouse monoclonal antibody. (UM800032; heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min)




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human colon tissue using anti-PSMA6 mouse monoclonal antibody. (UM800032; heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min)




Immunohistochemical staining of 3 cases of paraffin-embedded human endometrial carcinoma using anti-PSMA6 clone UMAB102 mouse monoclonal antibody (UM800032) at 1:200 with Polink2 Broad HRP DAB detection kit; heat-induced epitope retrieval with GBI Citrate pH6.0 HIER buffer using pressure chamber for 3 minutes at 110C. Cytoplasmic, membraneous and nuclear staining is seen in all 3 case of tumor cells.




Immunohistochemical staining of 3 cases of paraffin-embedded human lung using anti-PSMA6 clone UMAB102 mouse monoclonal antibody (UM800032) at 1:200 with Polink2 Broad HRP DAB detection kit; heat-induced epitope retrieval with GBI Citrate pH6.0 HIER buffer using pressure chamber for 3 minutes at 110C. Most of the staining is nuclear in positive cells of the lung.



Immunohistochemical staining of 2 cases of paraffin-embedded human placenta using anti-PSMA6 clone UMAB102 mouse monoclonal antibody (UM800032) at 1:200 with Polink2 Broad HRP DAB detection kit; heat-induced epitope retrieval with GBI Citrate pH6.0 HIER buffer using pressure chamber for 3 minutes at 110C. Staining is nuclear, cytoplasmic, and membraneous in trophoblasts cells of the placenta.



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-PSMA6 monoclonal antibody (Clone UMAB102).



Western blot of human tissue lysates (15ug) from 10 different tissues (1: Testis, 2: Omentum, 3: Uterus, 4: Breast, 5: Brain, 6: Liver, 7: Ovary, 8: Thyroid, 9: Colon, 10: Spleen). Dilution: 1:500.