

Product datasheet for **UM800089**

PD1 (PDCD1) Mouse Monoclonal Antibody [Clone ID: UMAB197]

Product data:

Product Type:	Primary Antibodies
Clone Name:	UMAB197
Applications:	IF, IHC, WB
Recommended Dilution:	IHC 1:200, WB 1:500, IF 1:100..
Reactivity:	Human, Mouse, Monkey, Dog, Rat, Rabbit
Host:	Mouse
Isotype:	IgG2a
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human PDCD1 (NP_005009) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.5~1.0 mg/ml (Lot Dependent)
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	29.2 kDa
Gene Name:	programmed cell death 1
Database Link:	NP_005009 Entrez Gene 18566 Mouse Entrez Gene 301626 Rat Entrez Gene 486213 Dog Entrez Gene 5133 Human Q15116



[View online »](#)

Background:

This gene encodes a cell surface membrane protein of the immunoglobulin superfamily. This protein is expressed in pro-B-cells and is thought to play a role in their differentiation. In mice, expression of this gene is induced in the thymus when anti-CD3 antibodies are injected and large numbers of thymocytes undergo apoptosis. Mice deficient for this gene bred on a BALB/c background developed dilated cardiomyopathy and died from congestive heart failure. These studies suggest that this gene product may also be important in T cell function and contribute to the prevention of autoimmune diseases. [provided by RefSeq, Jul

Synonyms:


CD279; hPD-1; hPD-I; hSLE1; PD-1; PD1; SLEB2

Protein Families:


Druggable Genome, Transmembrane

Protein Pathways:


Cell adhesion molecules (CAMs), T cell receptor signaling pathway

Product images:


Immunohistochemical staining of paraffin-embedded Human tonsil using anti-PDCD1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.0) at 120°C for 3min, UM800089) (1:200)




Immunohistochemical staining of paraffin-embedded VX2 tumor in rabbit liver using anti-PDCD1 mouse monoclonal antibody. (Heat-induced epitope retrieval by EDTA buffer (pH8.0) at 95-100°C for 30min, UM800089) (1:100)




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PDCD1 ([RC210364], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PDCD1 antibody (1:2000).




Western blot of cell lysates (35ug) from 9 different cell lines (1: HepG2, 2: HeLa, 3: SV-T2, 4: A549, 5: COS7, 6: Jurkat, 7: MDCK, 8: PC-12, 9: MCF7).




Western blot analysis of extracts (35ug) from PBMCs and PHA-stimulated-PBMCs by using anti-PDCD1 monoclonal antibody (1:250).



Western blot of human tissue lysates (15ug) from 10 different tissues (1: Testis, 2: Omentum, 3: Uterus, 4: Breast, 5: Brain, 6: Liver, 7: Ovary, 8: Thyroid, 9: Colon, 10: Spleen). Dilution: 1:500.



Immunofluorescent staining of PDCD1 ([RC210364])-stable-expression cells (left) labeling PDCD1 with mouse monoclonal antibody UM800089 (1:100, green) and nucleus with Hoechst33342 (blue). HEK293T cells serve as negative control (right).



Immunocytochemistry staining of PDCD1 ([RC210364])-stable-expression cells (left) labeling PDCD1 with mouse anti-PDCD1 monoclonal antibody UM800089 (1:900). The right is negative control.